Influence of Fire on Mercury Cycling in Boreal Forests

Randy Kolka (and many others)
Team Leader & Research Soil Scientist




Superior National Forest Research Forum February 22, 2012



Outline


- Background on Hg
 - Health Risks
 - Sources and Forms of Hg
 - Landscape Influences
- Fire Effects on Mercury Cycling
 - Fire Effects on Mercury Deposition
 - Fire Effects on Soil Mercury
 - Relationships between Forest Floor and Fish Mercury
- Summary

- Bioaccumulation in the Aquatic Food Chain
 - BAFs of ~ 1 million in Humans
 - New Research Terrestrial Bioaccumulation? Not really!



- **Health Consequences**
 - Attacks Neural Network and Nervous System
 - -Reproductive System
 - -Mercury Poisoning
 - Mad Hatters Disease
 - Iraq Poisoning
 - Minamata Disease

Not until the mid-1950's did people begin to notice a "strange disease". Victims were diagnosed as having a degeneration of their nervous systems. Numbness occurred in their limbs and lips. Their speech became slurred, and their vision constricted. Some people had serious brain damage, while others lapsed into unconsciousness or suffered from involuntary movements. Furthermore, some victims were thought to be crazy when they began to uncontrollably shout. People thought the cats were going insane when they witnessed "suicides" by the cats. Finally, birds were strangely dropping from the sky.



- Susceptible Groups
 - -Fetuses and Young Children
 - -Women of ChildbearingAge
 - Native Americans, Asians and Pacific Islanders



- Fish Consumption Advisories
 - Now in Every US state,Canadian Province andAcross Europe
 - -In MN All Surface Waters Have Warnings

WARNING

FISHERMAN


Fish taken from Cottage Grove Lake frequently contain elevated levels of mercury that can be harmful to health. It is believed that the source of mercury is normal geological mercury in the rocks and soil of this area.

The Oregon Health Division and Lane County Health Department advise you to limit your eating of these fish as follows:

- Pregnant women, nursing women and children up to six years of age should not consume any fish from this reservoir:
- Children older than six years and healthy adults should limit their consumption of fish from this reservoir to no more than one half pound (8 ounces) of fish from this reservoir per week.


If you have questions about this advisory please call Lane County Health Department at 687-3731







Background – Health Risks (e.g.)



Pregnant Women, Women who may become pregnant and Children under age 15

RIVER NAME	Species	Unrestricted	1 meal/week	1 meal/ month	Do not eat	Contaminants
St. Louis Fond du Lac Dam to Lake Superior	Crappie, Sunfish		All sizes			Mercury
	Perch		All sizes			Mercury
	Northern Pike			All sizes		Mercury
	Walleye			Shorter than 20 inches	Longer than 20 inches	Mercury & PCBs
	Smallmouth Bass, Catfish			All sizes		Mercury & PCBs


General Population

RIVER NAME	Species	Unrestricted	1 meal/week	1 meal/ month	Do not eat	Contaminants
St. Louis Fond du Lac	Crappie, Sunfish	All sizes				Mercury
Dam to Lake Superior	Perch		All sizes			Mercury
Superior	Northern Pike		All sizes			Mercury
	Walleye		Shorter than 20 inches	Longer than 20 inches		Mercury & PCBs
	Smallmouth Bass, Catfish		All sizes			Mercury & PCBs

Guidelines for pregnant women, women planning to become pregnant and children under age 15


Kind of fish you eat	How often can you eat it?"			
Fish caught in Minnesota:				
Sunfish, crappie, yellow perch, bullheads	1 meal a week			
Walleyes shorter than 20 inches, northern pike shorter than 30 inches, smallmouth bass, largemouth bass, channel catfish, flathead catfish, white sucker, drum, burbot, sauger, carp, lake trout, white bass, rock bass, whitefish, other species	———— 1 meal a month			
Walleyes longer than 20 inches, northern pike longer than 30 inches, muskellunge	— Do not eat.			
Commercial fish:				
 Shark, swordfish, tile fish, king mackerel 	—— Do not eat.			
Other commercial species, including canned tuna	See MDH's brochure, "An Expectant Mother's Guide to Eating Minnesota Fish."			
* These guidelines apply even if eating fish just during a vacation or for just one season.				

Guidelines for men, and for women not planning to become pregnant




Background - Sources of Hg

- Natural Sources of Hg
 - Not Many
 - Geologic Materials
 - Cinnebar
 - Volcanic
 - Carbon Deposits




Background – Sources of Hg

- Anthropogenic Sources of Hg
 - Power Plants
 - Other Fossil Fuel Combustion
 - Manufacturing






Background – Global Geographic Sources of Hg




Pacyna and Munthe, 2004

Background – Sources & Forms of Hg

- Atmospherically Derived
 - Precipitation + Dry Deposition
 - Local to Global Sources
- Important Forms
 - Gaseous (Elemental)
 - Ionic
 - Organic Methyl Hg (HARMFUL!)






Background – Landscape Influences


- Production of Methyl-Hg
 - Low Oxygen
 - Food (Carbon)
- WETLANDS
- LAKE/STREAM BOTTOM





Background – Landscape Influences

• WETLANDS



Fire Effects on Mercury Cycling

- Fire Effects on Mercury Deposition
- Fire Effects on Soil Mercury
- Relationship
 Between Forest
 Floor and Fish
 Mercury



Fire Effects on Hg Cycling

Hypothesis: Burning leads to greater Hg deposition and subsequent short-term elevated Hg in fish









Fire Effects on Hg Cycling

- Blowdown in BWCA of the Superior NF
- ~200,000 ha
- Five control and five burned watersheds
- Also had two wildfires during the study (Alpine and Cavity Lake)
- Measuring Hg in soil, precipitation, throughfall, lake water and fish



Fire Effects on Hg Deposition

- Fire leads to higher short-term deposition of Hg (both Total and Methyl), especially in conifer systems
- Increase is about 30-40% of annual deposition
- Can we pick up that pulse of Hg in the food chain?




Witt et al., 2009b


Fire Effects on Soil Hg (Short-term)


- Forest Floor
 - Areas with blowdown
 followed by salvage logging
 and then fire has lower Hg in
 forest floor
 - Here fire alone did not change Hg in forest floor (unlike most other studies)
 - Mineral Soils
 - No differences

B = Blowdown, S = Salvage Logging, F = Fire, C = Control, (Mitchell et al. 2012)



Fire Effects on Soil Hg (Long-term)






- Soil Hg lower in areas more recently burned
- Close relationship between %C and Hg

Relationship Between Forest Floor Hg and Fish Hg

As the % organic matter (%C) in the forest floor increases fish Hg increases



(Gabriel et al. 2012)

- Fire decreases Hg in forest floor and possibly A horizons
- Relationship between watershed soil OM and fish Hg
- Increased deposition the year following fire, but then lower Hg soils
- Perhaps a blip in fish Hg following fire but then probably long-term decreases because the watershed pool is smaller

Summary – Fire Effects on Hg Cycling

- Fire leads to increased Hg deposition in downwind areas following fire.
- Deposition can be ~30-40% of annual deposition.
- Possibly then, fire could lead to short-term Hg increases in fish.
- Fire lowers Hg in watershed soils.
- Positive relationship between watershed soil Hg and fish Hg.
- Theoretically then, fire should lead to long-term decreases in fish Hg Still working on it...
- Possible management recommendations include imposing a short-term
 2-5 year consumption advisory on lakes affected by fire.
- Following fire, good to mitigate soil erosion. Mercury is closely associated with carbon, so keeping O and A horizon material out surface water bodies is important.

Thank You!



Questions?